



08/01/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Assistant Commissioner for Patents
Washington, D.C. 20231

August 1, 2000

S I R:

Transmitted herewith for filing are the specification and claims of the patent application of:

John F. O'Connor, Galina I. Kovalevskaya, and Steven Birken for
Inventor(s)

METHODS FOR PREDICTING PREGNANCY OUTCOME IN A SUBJECT BY hCG ASSAY
Title of Invention

Also enclosed are:

7 sheet(s) of informal formal drawings.

Oath or declaration of Applicant(s). (unsigned)

A power of attorney (unsigned)

An assignment of the invention to _____

A Preliminary Amendment

A verified statement to establish small entity status under 37 C.F.R. §1.9 and §1.27.

The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT

	NUMBER FILED	=	NUMBER EXTRA*	X	RATE		FEE	
					SMALL ENTITY	OTHER ENTITY	SMALL ENTITY	OTHER ENTITY
Total Claims	24 -20	=	4	X	\$9.00	\$18.00	= \$	\$72.00
Independent Claims	9 -3	=	6	X	\$39.00	\$78.00	= \$	\$468.00
Multiple Dependent Claims Presented:	_____ Yes <input checked="" type="checkbox"/> No				\$130.00	\$260.00	= \$	\$ 0

*If the different in Col. 1 is less than zero, enter "0" in Col. 2

BASIC FEE	\$345.00	\$690.00
TOTAL FEE	\$	\$ 1230.00



Applicants: John F. O'Connor, et al.
U.S. Serial No.: Not Yet Known (continuation of PCT/US99/02289
filed 3 February 1999)

Filed: Herewith

Letter of Transmittal
Page 2

A check in the amount of \$ 1230.00 to cover the filing fee.

Please charge Deposit Account No. _____ in the amount of \$ _____.

The Commissioner is hereby authorized to charge any additional fees which may be required in connection with the following or credit any over-payment to Account No. 03-3125:

Filing fees under 37 C.F.R. §1.16.

Patent application processing fees under 37 C.F.R. §1.17.

The issue fee set in 37 C.F.R. §1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. §1.311(b).

Three copies of this sheet are enclosed.

A certified copy of previously filed foreign application No. _____ filed in _____ on _____ Applicant(s) hereby claim priority based upon this aforementioned foreign application under 35 U.S.C. §119.

Other (identify) Express Mail Certificate of Mailing No. EL628788540US; one loose set of figures 1-7B (7 Sheets)

RECEIVED

Respectfully submitted,



John P. White
Registration No. 28,678
Attorney for Applicants
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400



08/01/00

08-02-00

A

Dkt. 54205-A-PCT-US/JPW/GJC

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : John F. O'Connor, et al.

U.S. Serial No. : Not Yet Known (Continuation Application of PCT/US99/02289, filed 3 February 1999)

Filed : Herewith

For : METHODS FOR PREDICTING PREGNANCY OUTCOME IN A SUBJECT BY hCG ASSAY

1185 Avenue Of The Americas
 New York, New York 10036
 August 1, 2000

JC857 U.S. PTO
 09/630215

 08/01/00

Assistant Commissioner for Patents
 Washington, D.C. 20231
 Box: Patent Application

EXPRESS MAIL CERTIFICATE OF MAILING
FOR ABOVE-IDENTIFIED APPLICATION

"Express Mail" mailing label number: EL 628 788 540 US
 Date of Deposit: August 1, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. §1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents and Trademarks, Washington, D.C. 20231.

Printed Name: Samuel Khass

Respectfully submitted,

John P. White
 Reg. No. 28,678
 Attorney for Applicants
 Cooper & Dunham LLP
 1185 Ave of the Americas
 New York, New York 10036
 (212) 278-0400

09/01/00 08/01/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant : John F. O'Connor, et al.
U.S. Serial No. : Not Yet Known (Continuation Application
of PCT/US99/02289, filed 3 February
1999)
Filed : Herewith
For : METHODS FOR PREDICTING PREGNANCY
OUTCOME IN A SUBJECT BY hCG ASSAY

1185 Avenue Of The Americas
New York, New York 10036
August 1, 2000

Assistant Commissioner for Patents
Washington, D.C. 20231
Box: Patent Application

Sir:

**PRELIMINARY AMENDMENT TO THE ACCOMPANYING CONTINUATION
APPLICATION FILED UNDER 37 C.F.R. §1.53**

Applicants request that the following amendment be made in the
above-identified application:

In the Specification:

On page 1, after the title and before line 5, please delete the
paragraph and insert the following new sentence:

--This application is a continuation of PCT International
Application No. PCT/US99/02289, filed 3 February 1999,
designating the United States of America, which is claiming
the priority of U.S. Serial No. 09/017,976, filed February
3, 1998, the contents of which are hereby incorporated by
reference into the present application.--

RECEIVED SEP 15 2000

In the Claims:

Please cancel claims 4-5, 7, 9-13, 16-17, 19, 22-26, 29-30, 32-33, 35-37, 40-41, 45, 47-48, 50, 52, 54-55, and 57 without prejudice or disclaimer to applicants' right to pursue the subject matter of these claims in a future continuation or divisional application and amend claims 34, 43, and 56 pursuant to the provisions of 37 C.F.R. 1.121(b) by deleting the bracketed material and inserting the underlined material as follows:

- 34. (Amended) The diagnostic kit of claim 31[, 32 or 33] further comprising a control sample [(s)] wherein the control sample is from a normal pregnant subject [sample (s)].--
- 43. (Amended) The early pregnancy associated isoform of hCG recognized by the monoclonal antibody of claim [40] 38, wherein the monoclonal antibody is designated B152.
- 56. (Amended) The method of claim [48] 44, wherein the gestational trophoblast disease is choriocarcinoma or hydatidiform mole and the second antibody is B109.

REMARKS

This application is a continuation of PCT International Application No. PCT/US99/02289, filed 3 February 1999, designating the United States of America and claiming priority of U.S. Serial No. 09/017,976, filed February 3, 1998. Accordingly, the parent application, PCT International Application No. PCT/US99/02289, is pending today in the United

001050 "S F E E S S O

John F. O'Connor, et al.
U.S. Serial No.: Not Yet Known
(Continuation of PCT/US99/02289,
filed 3 February 1999)
Filed: Herewith
Page 3

States of America pursuant to 35 U.S.C. §363, and the subject continuation application is co-pending therewith in fulfillment of the provisions of 35 U.S.C. §120.

By this Preliminary Amendment, applicants have hereinabove amended the specification on page 1 to insert the continuation data. Applicants maintain that the amendments to the specification present no issue of new matter and are fully supported by the specification. Applicants have amended the specification to recite the continuing data for the above-identified application.

Claims 1-57 were pending in the subject application. By this Amendment applicants have canceled claims 4-5, 7, 9-13, 16-17, 19, 22-26, 29-30, 32-33, 35-37, 40-41, 45, 47-48, 50, 52, 54-55, and 57 without prejudice or disclaimer to applicants' right to pursue the subject matter of these claims in a future continuation or divisional application. Applicants have hereinabove amended claims 34, 43, and 56. Support for these amendments may be found inter alia on page 5, lines 13-19, page 11, lines 21-24, and page 12, lines 5-11. Accordingly, upon entry of this Amendment, claims 1-3, 6, 8, 14-15, 18, 20-21, 27-28, 31, 34, 38-39, 42-44, 46, 49, 51, 53, and 56 will be pending and under examination.

If a telephone interview would be of assistance in advancing prosecution of the subject application, applicants' undersigned attorney invites the Examiner to telephone at the number provided below.

00T000" S T 0 0 S 0 0 0

**Application
for
United States Letters Patent**

007080" 5720E560

To all whom it may concern:

Be it known that (we) John F. O'Connor, Galina I. Kovalevskaya, and
Steven Birken

have invented certain new and useful improvements in

METHODS FOR PREDICTING PREGNANCY OUTCOME IN A SUBJECT BY hCG ASSAY

of which the following is a full, clear and exact description.

METHODS FOR PREDICTING PREGNANCY OUTCOME
IN A SUBJECT BY hCG ASSAY

This application is a continuation-in-part application of U.S. Serial No. 09/017,976, filed February 3, 1998, the contents of which are hereby incorporated by reference into this application.

5

The invention disclosed herein was made with United States Government support under National Institutes of Health Grant Nos. NIEHS ES-07589 and HD 15454. Accordingly, the U.S. Government has certain rights in this invention.

10

Background of the Invention

Throughout this application, various publications are referenced by author and date. Full citations for these publications may be found listed alphabetically at the end of the specification immediately preceding the claims. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art.

15

20

Early pregnancy loss (EPL) is a widespread, but largely undiagnosed problem. In order to adequately diagnose and develop treatments for EPL it is essential to be able to detect and measure the rate of occurrence of EPL. This is critically important in epidemiological studies, some of which are related to exposures to known or suspected reproductive toxins in the workplace, in the environment or by personal use. These early pregnancy losses are often not recognized by women or physicians and are detected solely by the measurement of hCG in the urine at the time between implantation and expected menses. They are sometimes termed "chemical pregnancies" or "occult pregnancies." A landmark epidemiological study established that the incidence of

25

30

35

DOCTORED AT PAGE 360

EPL was 22% in a population of healthy women attempting to conceive (Wilcox, A. J., et al., 1988). This investigation employed a very sensitive (0.01 ng/ml hCG) assay which detected only the intact hCG molecule with the unique beta subunit carboxyterminal peptide present.

There are multiple likely causes for EPL and clinical spontaneous abortion including genetic abnormality, immunological dysfunction, untreated infection or other unknown physiological problems. In addition, losses may be caused by failure of human chorionic gonadotropin (hCG) to induce adequate response at its target, the corpus luteum. This could result from inadequate hormonal potency. "Nicking" of the beta subunit in the loop 2 region of the molecule, specifically between residues 44-49, can reduce biopotency of hCG. Cleaved peptide bonds in this area of the molecule also exhibit reduced biopotency and reduced immunochemical recognition by monoclonal antibodies directed to the heterodimeric hormone (Cole, L. A., et al., 1991a; Cole, L. A., et al., 1991b; Puisieux, A., et al., 1990; Nishimura, R., et al., 1988; Nishimura, R. T., et al., 1989). Nicked forms of hCG were examined as possibly more prevalent in EPL situations and, at least in part responsible, for early pregnancy loss. Unfortunately many of the reports claiming that substantial concentrations of nicked hCG are produced during pregnancy, losses or successful pregnancies, are not accurate due to faulty assumptions regarding assay specificity. Carbohydrate-modified hCG can also exhibit either reduced or enhanced biopotency. It is known that if the hCG has much reduced sialic acid content and its carbohydrate chains terminate in galactose, much hCG would be removed by the liver receptor for such altered glycoproteins (Braun, J. R., et al., 1996; Kawasaki, T. and G. Ashwell, 1996). The circulating life-time of asialo hCG is reduced and its *in vivo* potency is thereby

001080" 5120E950

low. Other carbohydrate changes also alter circulating half life; glycoproteins terminating in sulfate-N-acetyl galactosamine are also extracted by a specific liver receptor and have reduced circulating lifetime (Baenziger, J. U., 1994; Fiete, D., et al., 1991).

At least two factors affect increased potency of hCG. First, it is known that a larger Stoke's radius will decrease clearance through the kidney glomerulus which generally clears proteins above an effective size of 70,000 very slowly. The effective size of urinary-isolated hCG is just at this borderline reduced clearance size. Generally, extra sugar content makes the hydrated radius of glycoproteins larger. It has been shown that by adding the hCG beta COOH-terminal peptide to hFSH or hLH, their circulating life-times greatly increased (Fares, F. A. et al., 1992; Matzuk, M. M., 1990). This addition was thought mostly due to the carbohydrate content of that peptide rather than simply the extra polypeptide size. Second, increased negative charge of a protein will prolong its circulating time because of decreased renal clearance (Chmielewski, C. 1992, Quadri, K. H., et al., 1994; Maack, T., et al., 1985). This increased negative charge can be due to extra sialic acid or other negative groups, including sulfate such as is present on hLH and on the pituitary form of hCG (Birken, S., et al., 1996b). Changes which affect signal transduction at the receptor may also affect biopotency of hCG. It is known that deglycosylated hCG has much reduced receptor potency (Ravindranath, N., et al., 1992; Sairam, M. R., and L. G., Jiang, 1992; Browne, E. S., et al., 1990; Sairam, M. R., 1989; Sairam, M. R., et al., 1988). Carbohydrate reduced forms of hCG also have reduced signal transduction (Amano, J., et al., 1990; Bahl, O. P., et al., 1995; Moyle, W. R., 1975).

According to the present invention EPL or recurrent spontaneous abortion is not due to an abnormal hCG form that has reduced potency, such as nicked hCG. Instead, the present invention provides evidence that in successful outcome pregnancies women usually produce forms of hCG which are very highly potent in very early pregnancy; the standard urinary reference preparations of hCG are less potent forms of the hormone produced later in pregnancy. The increased potency could be caused by a combination of factors from circulating half-life to increased receptor affinity or signal transduction or all of the preceding. Since hCG is low very early in pregnancy, it is logical to find a more potent form of hCG on a molar basis to carry out its function until production levels rise as the trophoblastic cellular mass increases. The present invention describes molecular and immunological tools and methods including an antibody, B152, described herein which recognizes the highly potent early pregnancy associated molecular isoforms of hCG. The determination of blood and urine profiles for the B152 hCG isoforms throughout healthy pregnancies can delineate the pattern of isoforms in successful pregnancies. These isoforms can be measured by immunoassay alone, obviating the need to perform complex isoelectric focusing studies or other separation techniques. Additionally, the methods described herein are applicable to large numbers of samples.

00000000000000000000

Summary of the Invention

The present invention provides a method of predicting pregnancy outcome in a subject by determining the amount of an early pregnancy associated molecular isoform of hCG in a sample comprising: (a) contacting a sample with an antibody which specifically binds to the early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (b) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample; and (c) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (b) with the amount determined for temporally matched, normal pregnant subject(s) wherein the relative absence of the early pregnancy associated molecular isoform of hCG in the sample indicates a negative outcome of pregnancy for the subject.

The present invention further provides a method of predicting pregnancy outcome in a subject by determining the amount of an early pregnancy associated molecular isoform of hCG in a sample comprising: (a) capturing an antibody which specifically binds to the early pregnancy associated molecular isoform of hCG with a solid matrix under conditions permitting binding of the antibody with the solid matrix; (b) contacting the bound matrix with the sample under conditions permitting binding of the antigen present in the sample with the capturing antibody; (c) separating the bound matrix and the sample; (d) contacting the separated bound matrix with a detecting antibody which specifically binds to hCG under conditions permitting binding of antibody and antigen in the sample; (e) measuring the amount of bound antibody on the bound matrix, thereby determining the amount of early pregnancy associated molecular isoform

COPIES OF 30

of hCG in the sample; and (f) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (e) with the amount determined for temporally matched, normal pregnant subject(s), wherein amounts of the early pregnancy associated molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG in temporally matched pregnant samples indicates a positive outcome, amounts of early pregnancy associated molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG in the non-pregnant samples indicates a negative outcome of pregnancy for the subject.

In addition, the present invention provides a method for determining the amount of early pregnancy associated molecular isoforms of in a sample comprising: (a) contacting the sample with an antibody which specifically binds to an early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; and (b) determining the amount of complexes formed thereby determining the amount of early pregnancy associated molecular isoform of hCG in the sample.

Further, the present invention provides a diagnostic kit for determining the amount of early pregnancy associated hCG is a sample comprising: (a) an antibody which specifically binds to an early pregnancy associated molecular isoform; (b) a solid matrix to which the antibody is bound; and (c) reagents permitting the formation of a complex between the antibody and a sample.

The present invention additionally provides an antibody

which specifically binds to an early pregnancy associated molecular isoform of human chorionic gonadotropin.

5 Further, the present invention provides a method for
detecting non-trophoblast malignancy in a sample
comprising: (a) contacting a sample with an antibody
which specifically binds to the early pregnancy
associated molecular isoform of hCG under conditions
10 permitting formation of a complex between the antibody
and the early pregnancy associated molecular isoform of
hCG; (b) contacting the sample with a second antibody
which specifically binds to intact non-nicked hCG
without substantially cross-reacting with said antibody
15 under conditions permitting formation of a complex
between the antibody and the early pregnancy associated
molecular isoform of hCG; (c) measuring the amount of
complexes formed, thereby determining the amount of the
early pregnancy associated molecular isoform of hCG in
20 the sample; and (d) comparing the amount of early
pregnancy associated molecular isoform of hCG in the
sample determined in step (b) with the amount of early
pregnancy associated molecular isoform of hCG in the
sample determined in step (c), wherein a positive
25 detection of early pregnancy associated molecular
isoform detected in step (b) and a relative absence of
the early pregnancy associated molecular isoform of hCG
detected in step (c) indicates the presence of non-
trophoblast malignancy in the sample.

30

Finally, the present invention provides a method for
detecting gestational trophoblast disease in a sample
from a subject comprising (a) contacting a sample with
an antibody which specifically binds to the early
pregnancy associated molecular isoform of hCG under
35 conditions permitting formation of a complex between the
antibody and the early pregnancy associated molecular

isoform of hCG; (b) contacting the sample with a second antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (c) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample due to binding with the first antibody, and late pregnancy associated molecular isoform of hCG in the sample due to binding with the second antibody; (d) determining the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the subject; and (e) comparing the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) over time, wherein a continuing high ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) indicates the presence of gestational trophoblast disease in the subject.

5

10

15

20

25

007030 "STEP 00300

Brief Description of the Figures

Figure 1.

Bioassay for forms of hCG. This is data from recombinant CHO cells expressing the LH/CG receptor. The response factor is cAMP production. The x-axis is dose of one of four calibrated, pure hormones as described on graph legends. Expressed hCG has no nicks; choriocarcinoma hCG (C5) is 100% nicked; CR 127 was purified into a nick-free (intact) and nick-enriched fraction as shown.

Figure 2.

Incidence (Panel A) and expression level (Panel B) of hCG-related molecules in the positive samples for each of the analyses measured (In early normal pregnancy, n=214; EPL cycles, n=49; and negative cycles, n=297).

Figure 3.

Binding curves for three hCG types in the B152-B207* assay (upper panel) and the B109-B108* assay (lower panel).

Figure 4.

Ratio of hCG isoforms measured by the B152-B207* and B109-B108* assays in normal pregnancy urine (n=103) at different gestational ages. (Regression curve and 95% confidence intervals are shown, $r^2=0.79$). An inflection point in the curve occurs at approximately 29 weeks.

Figure 5.

Box plot of the B152/B109 ratio for pregnancy matched serum/urine at 5-6 weeks of gestational age (n=12); or at 36-39 weeks of gestational age (n=11) and in JAR cell supernatant. Box extends to the 25th and 75th percentile. The upper and lower symbols indicate the 90th and 10th percentile respectively. A solid line inside the box marks the value of the 50th percentile.

Figure 6.

Ratio of hCG isoforms measured by the B152-B207* and B109-B108* assays in the urine of IVF patients (n=65). (Regression curve and 95% confidence intervals are shown, $r^2=0.59$).

5

Figure 7.

Immunoassay profiles of fractions from Superose 12 column chromatography of a pooled urine concentrate from pregnant women.

10

001030 " 57205960

5 said antibody under conditions permitting formation of
a complex between the antibody and the early pregnancy
associated molecular isoform of hCG. In an embodiment
of this invention, the second detection antibody is
10 B207. According to another embodiment of this invention,
step (a) further comprises a second assay antibody B109
which specifically binds to intact non-nicked hCG
without substantially cross-reacting with said antibody
under conditions permitting formation of a complex
15 between the antibody and the early pregnancy associated
molecular isoform of hCG. In an embodiment of this
invention, the detection antibody is B108. In an
embodiment of this invention, step (c) comprises
comparing the amount of the early pregnancy associated
20 molecular isoform of hCG determined in step (b) for
B152-B207 assay with the amount determined in step (b)
for the B109-B108 assay wherein a high ratio of amounts
determined for said antibody relative to the second
antibody indicates a positive outcome of pregnancy for
the subject, a low ratio indicates a negative outcome of
pregnancy for the subject.

In yet another embodiment of this invention, step (c)
25 comprises comparing the amount early pregnancy
associated molecular isoform of hCG in the sample
determined in step (b) with the amount determined for
temporally matched, normal pregnant subject(s), wherein
amounts of the early pregnancy associated molecular
30 isoform of hCG in the sample similar to amounts of early
pregnancy associated molecular isoform of hCG in
temporally matched pregnant samples indicates a positive
outcome, amounts of early pregnancy associated molecular
isoform of hCG in the sample similar to amounts of early
35 pregnancy associated molecular isoform of hCG in the
non-pregnant samples indicates a negative outcome of
pregnancy for the subject.

COPIED FROM STREETS

embodiment of this invention, the capturing antibody is B152. In one embodiment of this invention, the detecting antibody is B207. In an embodiment of this invention, step (a) further comprises a second capturing antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG. According to an embodiment of this invention, the second capturing antibody is B109 and the second detection antibody is B108. In an embodiment of this invention, step (d) further comprises a second detecting antibody which specifically binds to hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG. In an embodiment of this invention, step (f) comprises comparing the amount of the early pregnancy associated molecular isoform of hCG determined in step (e) for said antibody with the amount determined in step (b) for the second antibody, wherein a high ratio of amounts determined for said antibody relative to the second antibody indicates a positive outcome of pregnancy for the subject, a low ratio indicates a negative outcome of pregnancy for the subject.

According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least two consecutive days. In an embodiment of this invention, the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least two consecutive days. In one embodiment of this invention, the antibody is labeled with a detectable marker. In an embodiment of this invention,

001050 "SECRET" 030100

According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least one or may be two or three consecutive days. In an embodiment of this invention, the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least one, or may be two or three consecutive days. In one embodiment of this invention, the antibody is labeled with a detectable marker. In an embodiment of this invention, the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive isotope is I^{125} .

The present invention additionally provides an antibody which specifically binds to an early pregnancy associated molecular isoform of human chorionic gonadotropin.

In an embodiment of this invention, the antibody specifically binds to a region of the early pregnancy associated molecular isoform of human chorionic gonadotropin comprising a carbohydrate moiety.

According to one embodiment of this invention, the monoclonal antibody is B152. In an embodiment of this invention, a hybridoma cell (ATCC Accession No. HB-12467) is provided capable of producing monoclonal antibody B152. Another embodiment of this invention is the early pregnancy associated molecular isoform of hCG recognized by the B152 monoclonal antibody.

Further, the present invention provides a method for detecting non-trophoblast malignancy in a sample comprising: (a) contacting a sample with an antibody which specifically binds to the early pregnancy associated molecular isoform of hCG under conditions

099001-510000

5 permitting formation of a complex between the antibody
and the early pregnancy associated molecular isoform of
hCG; (b) contacting the sample with a second antibody
which specifically binds to intact non-nicked hCG
under conditions permitting formation of a complex
between the antibody and the early pregnancy associated
10 molecular isoform of hCG; (c) measuring the amount of
complexes formed, thereby determining the amount of the
early pregnancy associated molecular isoform of hCG in
the sample; and (d) comparing the amount of early
pregnancy associated molecular isoform of hCG in the
15 sample determined in step (b) with the amount of early
pregnancy associated molecular isoform of hCG in the
sample determined in step (c), wherein a positive
detection of early pregnancy associated molecular
isoform detected in step (b) and a relative absence of
the early pregnancy associated molecular isoform of hCG
detected in step (c) indicates the presence of non-
20 trophoblast malignancy in the sample.

According to an embodiment of this invention, the
antibody is B152 or B109. In an embodiment of this
invention, the detection antibody is B207 for B152
25 assay, B108 for B109 assay. In an embodiment of this
invention, the non-trophoblast malignancy is ovarian
malignancy or prostate malignancy.

30 According to an embodiment of this invention, the sample
is a urinary sample or a blood sample. In one
embodiment of this invention, the sample is an aggregate
sample taken from at least two consecutive days. In an
embodiment of this invention, the sample is a spot urine
sample, a first morning void urine sample, or an
35 aggregate sample of the first morning void urine samples
for at least two consecutive days. In one embodiment of
this invention, the antibody is labeled with a

00000000000000000000

detectable marker. In an embodiment of this invention, the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive isotope is I¹²⁵.

5

Finally, the present invention provides a method for detecting gestational trophoblast disease in a sample from a subject comprising (a) contacting a sample with an antibody which specifically binds to the early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (b) contacting the sample with a second antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (c) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample due to binding with the first antibody, and late pregnancy associated molecular isoform of hCG in the sample due to binding with the second antibody; (d) determining the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the subject; and (e) comparing the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) over time, wherein a continuing high ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) indicates the presence of gestational trophoblast disease in the subject.

10

15

20

25

30

35

In an embodiment of this invention, the antibody is B152

007030 "SECRET" 000000

or B109. In another embodiment of this invention, the detection antibody is B108 for B109 assay, B207 for B152 assay. In an embodiment of the present invention, the gestational trophoblast disease is choriocarcinoma or hydatidiform mole.

According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least two consecutive days. In an embodiment of this invention, the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least two consecutive days. In one embodiment of this invention, the detection antibody B207 or B108 is labeled with a detectable marker. In an embodiment of this invention, the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive isotope is I¹²⁵.

As described herein below, unexpected isoforms of hCG are produced during normal early pregnancy. Using an *in vitro* bioassay, it appears that these isoforms have enhanced potency for signal transduction. These isoforms can be measured using the novel sensitive, immunoassay described herein. This can help predict pregnancy outcome where one cause of early pregnancy loss is failure to produce the isoform of hCG of higher potency produced by successful pregnancies. This enables physicians to intervene to sustain a failing pregnancy. Identification of the nature of the hCG isoform required might provide the proper reagent needed to sustain pregnancy.

New antibodies for measurement of nicked forms of hCG described herein below were developed based on the hypothesis that forms of hCG, which have greatly reduced

001030" 5720E950

bioactivity, contribute to early pregnancy loss (EPL), due at least in part to diminished biopotency. Evidence was found that the hCG that appears in EPL patients displays reduced biological activity. However, it was

5 determined that the cause of the reduced bioactivity is not the presence of nicked hCG in EPL patients. Instead, the hypothesis is that patients that carry pregnancies forward produce an isoform of hCG with enhanced bioactivity. The instant invention describes

10 a unique immunochemical assay to measure this unexpected and previously un-characterized isoform of early pregnancy hCG directly in clinical samples of blood and urine. One of the antibodies developed reacted against a nicked form of hCG isolated from a choriocarcinoma

15 patient, was not specific for a nicked form of hCG but appeared to discriminate among carbohydrate variants of hCG. This antibody, designated B152, appears to preferentially bind hCG forms from choriocarcinoma patients. In studying the content of hCG isoforms

20 during pregnancy, the unique and unexpected observation was made that B152 in the first four weeks of pregnancy measured much higher quantities of an isoform of hCG as compared to the standard hCG isoforms measured by the usual heterodimeric hCG assays exemplified by a

25 previously described B109 based assay. In fact, in early pregnancy (days 9,10,11 postovulation) B152 measured as much as 20-fold more hCG, than did another monoclonal antibody, B109. Later in pregnancy, the B152 isoform declines and is lower in third trimester

30 pregnancy urine than the standard isoforms measured by B109. A further striking observation was that in very early pregnancy, a high B152/B109 ratio correlates with a successful pregnancy outcome while a low ratio correlated with pregnancy loss. This discovery is

35 important as the potentially overlooked isoforms of hCG described herein during pregnancy may be predictors of successful pregnancy outcome. Such an assay has wide

000000000000000000000000

medical applications and provides a clinician with opportunity to intervene very early in pregnancy if the assay indicated that the pregnancy appeared troubled.

5 An antibody, designated B152, produced by the hybridoma cell accorded ATCC Accession number HB-12467 generated against a nicked form of hCG isolated from a choriocarcinoma patient, but not specific for nicked isoform hCG is able to discriminate among carbohydrate
10 variants of hCG. B152 is specific for an early pregnancy associated molecular isoform of hCG. which in the first four weeks of pregnancy is measured at much higher quantities than the hCG standard isoforms measured by the usual heterodimeric hCG assays
15 exemplified by a previously described B109 based assay. Later in pregnancy, the B152 isoform declines and is lower in third trimester pregnancy urine than the standard isoforms measured by B109.

20 This invention is illustrated in the Experimental Details section which follows. These sections are set forth to aid in an understanding of the invention but are not intended to, and should not be construed to, limit in any way the invention as set forth in the
25 claims which follow thereafter.

000000 " 512000000

EXPERIMENTAL DETAILS

Example 1: Analysis of Molecular Isoforms of hCG in
Early Pregnancy and Early Pregnancy Loss

5

Introduction

Almost all investigations of the incidence of early pregnancy loss (EPL), either in normal populations or in populations at risk as a consequence of exposure to putative reproductive toxins (Hakim, R. B., et al., 1995; Lasley, B. L., et al., 1995) use assays for heterodimeric, non-nicked hCG or combination assays which additionally include free beta subunit and beta core fragment of hCG. One concern about the forms of hCG to include in the measurement in EPL was heightened with respect to the nicking phenomenon described above. Because nicked hCG molecules are not measured by the antibodies employed in most EPL studies, the incidence of EPL is presumably underestimated by an amount proportional to the extent of nicking in the urinary molecule. Another concern of significant importance was a determination of the nature of the "hCG like" immunoreactivity in the urine in the periovulatory surge of the menstrual cycle (O'Connor J., et al., 1995). Recent reports have confirmed the existence of and documented the structure of a sulfated form of hCG produced in the pituitary (Birken, S., et al., 1996b). There is a pulsatile secretion of hCG in both men and non-pregnant women. (Odell, W. D.; Griffin, J., 1989 and Odell, W. D.; Griffin, J., 1987). The presence of a non-pregnancy associated form of sulfated hCG of pituitary origin, peaking at ovulation and perhaps persisting into the luteal phase, could potentially interfere with the accurate estimation of EPL.

35

Unappreciated isoforms of hCG in blood and urine very early in pregnancy may be more potent *in vivo* than the forms of hCG produced later in pregnancy. The absence

001030" ST00E960

of such isoforms may be one cause of early pregnancy loss. A sensitive and specific immunoassay system was designed and made to measure unique early pregnancy associated molecular isoforms (EPMI) of hCG. These isoforms, likely to differ by carbohydrate composition, are predictive of a successful pregnancy outcome. When these early pregnancy associated molecular isoforms of hCG are absent or present in low concentration, the pregnancy may be lost very early and be observed as only a "chemical" pregnancy. These hCG isoforms may resemble the forms of hCG produced in some choriocarcinoma patients from which the immunogen used to produce monoclonal antibody B152 was derived as described herein below. The isoforms resemble those from trophoblastic disease not in terms of nicking or intact peptide chains but likely in carbohydrate content. The present invention describes that the molar ratio of B152 to B109 epitopes are predictive of a successful pregnancy or a loss. Three categories of pregnant patients were analyzed: (a) normal pregnant women, (b) women who experience recurrent abortions, (c) women undergoing embryo implantation.

It is possible to determine the hCG isoforms present in the blood and urine of women who have a history of recurrent spontaneous abortion and a similar analysis of women undergoing embryo implantation. The combined EPL and spontaneous abortion rate in healthy populations is 31%. Subjects who experience three consecutive recurrent spontaneous abortions have a 32% risk of sustaining another (Hill, J. A.; Anderson, D. J., 1990). In in vitro fertilization IVF pregnancy, the loss rate is 70% with non-donor sperm and 50% when donor sperm is used. Delineation of pregnancies with a negative outcome from pregnancies with a positive outcome can be based on differences in the concentrations of EPMI hCG isoforms (i.e. as differences in the B152/B109 ratio in

001000 " 57 00000

patients). In addition, specimens from gestational trophoblastic disease (GTD) can be used to discriminate between GTD and normal pregnancy.

Results

5 In vitro bioassay for hLH/hCG

An hCG bioassay was constructed employing CHO cells expressing functional human LH/CG receptor. Figure 1 illustrates the differences *in vitro* in biological activity between nicked and non-nicked hCG as measured by this assay. This system, has been used to evaluate the activity of pituitary and placental hCG (Birken, S., et al., 1996b). Preparations of hCG were tested for nicked and non-nicked molecular isoforms of hCG in a second recombinant bioassay system (Ho, H-H., et al., 10 15 1997). Similar results were obtained in both systems.

Normal pregnancy values compared with EPL values.

Figure 2 indicated that nicked hCG is not a significant molar constituent of either early pregnancy or EPL. Data indicated that biological activity is not correlated with nicked hCG, but is instead ascribed to a form of hCG recognized by the B152 monoclonal antibody -- an early pregnancy associated molecular isoform of hCG (EPMI hCG). It has been established that there is diminished hCG bioactivity associated with EPL as compared to early normal pregnancy (Ho, H-H., et al., 20 25 1997). Thus, diminished hCG biological activity is a factor in EPL as a consequence of a heretofore unappreciated isoform of hCG - an early pregnancy associated molecular isoform of hCG. 30

hCG Urinary Analytes. Metabolites of hCG and hLH were studied in a variety of states (Birken, S., et al., 1996a). One study indicated a 31% pregnancy loss (Zinaman, MJ, et al., 1996) while another indicated a 35 17.4% rate of early pregnancy loss based on hCG assays (Ellish, N. J., et al., 1996). It is known that hCG and

001030" 51200000

hCG beta core can be readily transferred from the uterus
to the circulation even in the absence of implantation
(Chang, P. L., 1997). The molecular spectrum of hCG
urinary analytes in EPL cycles, normal conceptive cycles
5 and non-conceptive cycles has been evaluated. The study
design and demographics of the investigation have been
described (Elish, N. J., et al., 1996).

Briefly, three urine specimens per cycle, corresponding
10 to days 9,10, 11, post calculated day of ovulation were
collected and analyzed in a screening assay (the
"combo") which simultaneously detects intact, non-nicked
hCG, hCG free beta subunit, and hCG beta core fragment.
Individual determinations for each of these analytes, as
15 well as for nicked hCG, and the form of intact hCG
detected by monoclonal antibody B152 (EPMI hCG) were
performed on these specimens. In addition, since the
concentration of luteal phase hLH urinary analytes is a
concern because of cross-reaction in hCG assays, levels
20 of intact hLH, hLH free beta subunit and hLH beta core
fragment were determined in the normal pregnancy cycles
and the non-conceptive cycles. Table I summarizes the
characteristics of immunometric assays employed.

25

TABLE 1. Assay format and specificity

<i>Assay format</i>	<i>Primary analyte</i>	<i>% cross-reactivity with related analytes</i>	<i>Intra-assay cv, %</i>	<i>Inter-assay cv, %</i>
B109-B108*	intact non-nicked hCG	<1% ^b	6	12
B201-C104*	hCG free beta subunit (non-nicked+nicked)	1% hCG; 10% hCG nicked (pregnancy); <1% ^b	6	12
B210-B108*	hCG beta core fragment	2% hLH beta core fragment; <1% ^b	5	7
B151-B207*	hCG nicked	10% hCG nicked free beta subunit; 12% hCG non-nicked; 2% hCG free beta subunit; 2% hLH; 5% hLH free beta subunit; <1% ^b	5	15
B152-B207*	choriocarcinoma hCG (C5) and choriocarcinoma hCG free beta subunit	100% hCG nicked (C5); 190% hCG free beta nicked (from C5); 10% hCG nicked (pregnancy); 5% hCG free beta nicked (pregnancy); 7% hCG (pregnancy); 6% hCG free beta subunit; <1% ^b	6	13
B406-A201*	hLH	<1% ^a	4	10
B505-B503*	hLH beta core fragment	<1% ^a	9	9
B408-B409*	hLH free beta subunit	29% hLH; <1% ^a	7	11

^a-(if not indicated) hLH, free beta hLH, hLH beta core fragment, hCG, free beta hCG, hCG beta core fragment;

^b-(if not indicated) the same as ^(a) plus nicked hCG and nicked free beta hCG (pregnancy).

D:\Data\57202550

The results indicate that nicked hCG does not constitute a significant mole fraction of urinary hCG immunoreactivity in either EPL or early normal pregnancy. In addition, there is a substantial excretion of hCG free beta subunit in some subjects in both pregnancy and EPL. Further, both EPL and normal pregnancy cycles variably express all of the measured analytes. Although both the incidence and level of expression are different between EPL's and normal pregnancy, there is no hCG related analyte unique to either state. There was, however, a clear difference between the hLH associated analytes in the control population (non-conceptive cycles) and the normal pregnancy group. Virtually all of the non-pregnancy cycles expressed hLH free beta subunit and hLH beta core fragment while only a third of the conceptive cycles had detectable levels of either analyte. Intact hLH proved to be a minor constituent of the hLH profile in both groups.

These findings demonstrate both the necessity of measuring hCG beta core fragment in the detection of EPL, and also of making sure that the hCG beta core assay does not cross-react with beta core hLH, which is demonstrated to be present in that part of the luteal phase where EPL measurements are performed. The data is summarized in Figure 2.

Statistical analysis was performed after transformation of analyte values to mole fractions so as to produce a more useful analysis due to the wide excursion of hCG analyte values among groups. The mole fraction data were evaluated by discriminant analysis and by a mixed effects model incorporating LMP. The discriminant analysis was performed both with and without "outliers" (defined as values greater than two standard deviation from the mean) removed. Both approaches produced

007020"57020350

similar results.

5 A quadratic discriminant analysis based on a cross-validation method in order to minimize bias correctly classified 91% of the normal pregnancy subjects and 80% of the EPL subjects.

10 The mixed effects analysis, testing for interactions between mole fraction of analyte and time since LMP found no significant time or group (EPL vs. normal) effects in the intact hCG assay. In the free beta subunit of hCG assay, there is a significant group effect but no time trend. In both the hCG beta core fragment measurement and the B152 measurement, both the hormone levels and the time trend from LMP were significantly different between the EPL and pregnancy groups. This study produced several important findings. It defined the spectrum of analytes which in both early pregnancy and EPL, thereby resolving the issue of which hCG analytes to measure in epidemiological studies in which EPL is the end point determination. More importantly, it illustrated for the first time that there are significant differences both in the pattern of analytes and the time course of their appearance between early normal pregnancy and EPL. This observation facilitates very early prediction of a distressed pregnancy by urinary hCG measurements at a time which would permit therapeutic intervention.

25
30 Immunoreactivity of different forms of hCG in the two IRMA's (B152-B207 and B109-B108)

35 The relative binding of three different forms of hCG (urinary hCG, pituitary hCG and choriocarcinoma hCG C5) has been characterized in the two hCG assays (Figure 3). Urinary non-nicked hCG and pituitary hCG are recognized nearly equally well by the two IRMA's, while C5

001000 51202350

recognition is quite different. The B152-B207* assay is more sensitive to C5, which is to be expected because B152 antibody was developed and selected on the basis of higher affinity to C5. Urinary non-nicked hCG is purified from the CR127 preparation of pooled normal pregnancy hCG. Conversely C5 is recognized with lower affinity by the B109-B108* assay, which has primary specificity for the hCG isoforms of later pregnancy.

We have developed a method to directly profile changes of hCG isoforms in serum or urine throughout pregnancy. Two IRMAs for hCG are employed, each based on monoclonal antibodies to different hCG epitopes. The B109-B108* assay is a commonly used intact hCG assay to the heterodimeric-dependent epitope. A new assay, B152-B207*, is most likely sensitive to the carbohydrate portion of hCG carboxyterminal peptide. The same standard non-nicked hCG was used in both assays. Non-nicked hCG was employed since the B109 assay reacts poorly with nicked forms of hCG while the B152 assay does not discriminate between nicked and non-nicked forms of the hormone. The B152 assay detected with greatly enhanced sensitivity hCG isoforms which appear earlier in pregnancy than isoforms measured by the B109 assay (O'Connor et al. 1998). Prior to development of the new immunometric assay system described in this report, it was not possible to readily discern the changes in hCG isoforms from very early pregnancy to mid pregnancy. The only available procedure for examining these changes was isoelectric focusing of every patient specimen followed by immunoassay of every focused fraction (Berger et al. 1993; Ulloa-Aguirre et al. 1990). The IEF pattern reflects the heterogeneity of the charged sugar, sialic acid which varies with the multi-antennary structures of the carbohydrate moieties in which sialic acid is the terminal sugar. Although we do not yet know the precise nature of the isoform epitopes

007030 STAGE 560

being measured, the evidence for carbohydrate discrimination is based upon the hyperglycosylated structure of the immunogen, C5, used to develop the B152 monoclonal antibody and the antibody's reactivity with the hCG isoforms found in the JAR choriocarcinoma cell line. C5 hCG was isolated from a choriocarcinoma patient and has been thoroughly characterized as to its protein and carbohydrate content and structure (Elliott et al. 1997). It has been shown that C5 (and hCG from other choriocarcinoma subjects) differ in the protein moiety mainly by the presence of an increased number of nicked sites and by increased glycosylation relative to the hCG of normal pregnancy. In comparison with the hCG of normal pregnancy, choriocarcinoma derived hCG has increased fucosylation of the N-linked biantennary oligosaccharides in the beta subunit. In addition, the O-linked oligosaccharides in preparation C5 (a form of hCG produced from a single patient with choriocarcinoma) has a 100% tetrasaccharide core on the COOH-terminal region of the beta subunit. Normal mid pregnancy hCG has only 10-20% of this structure (Elliott et al. 1997). These observations, plus our own determination that the hCG synthesized by the JAR choriocarcinoma cell line provides a B152/B109 isoform ratio similar to that observed in early pregnancy, leads us to the conclusion that in very early pregnancy, the developing trophoblast secretes an isoform of hCG which resembles that produced in choriocarcinoma.

We have also tested recognition of pituitary hCG since its N-Asn carbohydrates differ somewhat from those of placental hCG, bearing a closer resemblance to those of hLH which have both sialic acid and sulfate groups (Birken et al. 1996). The carbohydrate structure of the b COOH-terminal portion of pituitary hCG is not yet known. Since B152 did not recognize any substantial differences between pituitary and placental hCG (Figure

001080 51202960

Statistical analysis involved fitting the log transformed ratio data to second and third order polynomial regression models. Since the third order term was not significant (likelihood ratio $\chi^2(1)=1.32$,
5 $P=0.25$), the second order model was used ($r^2=0.793$). The log B152/B109 ratio reached an inflection point at LMP=29 weeks, based on this model.

The B152-B207* values reflect a measurement of the B152
10 isoform in terms of later pregnancy hCG equivalents, not in absolute quantities. It must be emphasized that the "absolute" concentrations measure in the B152 assay cannot be compared with the results of the B109 assay on an equimolar basis since the potency of the
15 hyperglycosylated isoform is much higher in the B152 assay vis-à-vis the standard, i.e. normal later first trimester pregnancy hCG. The actual molar values of this isoform are on the order of tenfold less than those recorded in the assay. For this reason we have chosen
20 not to analyze absolute molar quantities of the two analytes but only the ratio of the two measurements.

Even in normal pregnancy, the hCG values obtained vary widely according to the characteristics of the
25 immunological reagents employed (Cole and Kardana, 1992; Cole et al. 1993). We hypothesize that the two assays described in this report primarily detect hCG isoforms at opposing ends of this spectrum, each primarily recognizing a subset of closely related molecules in the
30 continuum of early to later pregnancy hCG molecular forms.

We have retained the use of normal pregnancy hCG as the
35 standard in B152-B207* assay, despite its decreased affinity in this antibody configuration. The reasons for this include the limited and unrenewable supply of C5 (which was isolated from the urine of a single

007080" 57222560

patient) and the variability in data which would result from investigations using different standards. The consequences of this choice are that the early pregnancy hCG isoforms have markedly increased immunopotency over that of normal pregnancy and hence their molar quantities are overestimated in this assay. We use this difference in affinity to our advantage by employing a ratio of the molar results of two assays (B152 and B109). Either assay taken alone obscures this change due to the wide excursion of hCG values which occur in normal pregnancy.

Others have documented progressive changes in hCG isoforms throughout pregnancy. Skarulis et al. found that the fucose content of both intact hCG and also its free beta subunit increased as pregnancy progressed (Skarulis et al. 1992). Diaz-Cueto et al. investigating the isoelectric focusing pattern of circulating hCG throughout pregnancy, found that in early pregnancy, more than 80% of the hCG isoforms were acidic. This fraction decreased to less than half (47%) late in the third trimester (Diaz-Cueto et al. 1996). In contrast, Wide and Hobson found that the hCG of early pregnancy was more "choriocarcinoma-like" by virtue of its greater biological activity than the hCG of normal pregnancy (Wide and Hobson, 1987). Fein et al., in a study which employed gel filtration determined that first trimester hCG was a larger size than that of the third trimester. Treatment with exoglycosidases eliminated the size differential, indicating that the first trimester hCG was more highly glycosylated (Fein et al. 1980).

007050 "STAGE 950

The B152/B109 ratio in matched serum/urine samples in the first and third trimesters of pregnancy compared with hCG from JAR cells.

5 The B152/B109 ratio in serum is analogous to that found in matched urine samples and undergoes a similar change as pregnancy progresses (Figure 5). The B152/B109 ratio in the cell supernatant from JAR cells (a choriocarcinoma derived cell line) was similar to that
10 of early pregnancy.

The B152/B109 ratios of both serum and urine hCG concentrations are significantly higher in the first trimester as compared to the third trimester of normal
15 pregnancies (Table 2). Significant differences between serum and urine hCG concentration ratios as well as log transformed ratios in early (5-6 weeks) and late (36-39 weeks) gestation were evaluated by paired t-tests (Table 3). In both the first and third trimesters, urinary
20 B152/B109 ratios were significantly higher than serum ratios, indicating that there was a preferential clearance of the B152-recognized isoform into urine, regardless of the relative concentrations of the two isoforms.

25

TABLE 2.

Analysis of the B152/B109 ratio in serum and in urine in the first vs third trimesters of pregnancy.

Measure	T-test (df)	P
Serum, ratio B152/B109	t (11)=6.65	0.0001
Serum, log(ratioB152/B109)	t (23)=21.61	0.0000
Urine, ratio B152/B109	t (11)=4.64	0.0007
Urine, log(ratioB152/B109)	t (15.7)=16.85	0.0001

TABLE 3.

Analysis of the B152/B109 ratio in serum vs urine in the first and third trimesters of pregnancy.

Gestational age	Measure	Paired-t (df)	P
5-6 weeks	Ratio B152/B109	t (11)=3.25	0.0077
	Log(ratioB152/B109)	t (11)=6.25	0.0001
36-39 weeks	Ratio B152/B109	t (10)=5.47	0.0003
	Log(ratioB152/B109)	t (10)=7.14	0.0001

The B152/B109 ratio in urine samples from IVF patients

In urine samples from IVF patients (1-4 weeks post embryo transfer - ET) the B152/B109 ratio was again
5 between 2-8 and decreased as pregnancy progressed (Figure 6), similar to that observed in natural conceptions. The effect of pregnancy duration with respect to outcome variables could best be represented by a linear or quadratic function. ANCOVA models
10 including the second order week were fitted to the general equation: Outcome= (effect of time post ET)+(effect of diagnosis). After an appropriate ANCOVA model was determined, the least square means (adjusted for week post ET effect) were compared among the normal pregnancy, ectopic pregnancy and spontaneous abortion
15 populations (Table 4). The log transformed values of both B109-B108* and B152-B207* measured hCG forms discriminated both ectopic pregnancy and spontaneous abortions from normal pregnancy (P=0.0001). The ratio of
20 the log transformed values discriminated abortion from normal pregnancy (P=0.016). However, neither the ratio of B152/B109 nor the log of that ratio discriminated either of the pregnancy disorders from normal pregnancy.

007080 "SFEES60

TABLE 4.

IVF patients: analysis of covariance of hCG isoforms among normal pregnancy (np), ectopic pregnancy and spontaneous abortion as a function of gestational age.

Outcome	^c Adjusted R ²	^e -F	P	^d -Pairwise Difference
^a Log (ratio B152/B109)	0.51	0.89	0.41	none
^a Log (B109-B108*)	0.56	21.33	0.0001	np vs abortion & ectopic
^b Log (B152)/log (B109)	0.45	4.34	0.016	np vs abortion
^b Log (B152-B207*)	0.50	26.94	0.0001	np vs abortion & ectopic

- a- ANCOVA model with 2nd order polynomial coefficient (or parameter).
- b- ANCOVA model with only 1st order (linear) coefficient.
- c- Adjusted R² is a R² adjusting number of coefficients on the ANCOVA model so that comparisons of two R² with different ANOVA models with different number of coefficients are meaningful.
- d- "Pairwise difference" is based on t-test comparing the least-square means of outcome variables (after adjusting effect of week ET).
- e- Degree of freedom (df1, df2) for F-test are (2,82) for a model with only linear coefficient and (2,81) for a model with both linear and 2nd order coefficient.

HCG analysis of trophoblastic disease samples

Trophoblast disease serum (17 samples) and urine (28 samples) were obtained from patients post therapy and hence contained low hCG levels. Due to limited amounts of sample all of these specimens were run at a 1:10 initial dilutions. HCG levels in serum were low. The highest hCG concentration in serum was 202 fmol/ml in

00T080"9T20E450

the B152-B207* assay, with a corresponding value of 148 fmol/ml in the B109-B108* determination. Six of seventeen samples in serum had detectable levels, with 4/6 having a higher value in the B152-B207* assay. Of the 15/28 positive urine samples however, 14/15 had higher levels in the B152-B207* assay than in the B109-B108* assay, with the highest hCG value being 20000 fmol/ml in the B152-B207* assay and 18715 fmol/ml in the corresponding B109-B108* assay. Due to the small sample size, no statistical treatment was performed on this data, but even in these post-treatment patients the B152/B109 ratio was ≥ 1 , which corresponds to the early pregnancy hCG isoform ratio.

The specimen limitations discussed above precludes our reaching any definitive conclusion on the analysis of trophoblastic disease samples. However it appears as might be anticipated that the B152 assay is more sensitive than B109 assay in detecting hCG immunoreactivity in the blood and in the urine of trophoblastic disease patients, even after treatment.

Chromatography of First Week of Gestation Pregnancy Pool. In order to determine whether the B152-B207* assay recognized other forms of hCG associated immunoreactivity in addition to the intact hCG molecule, specimens were pooled. FPLC on tandem Superose 12 columns followed by immunoassay of the fractions for all of the characterized forms of hCG revealed that only the intact hCG molecule (or hCG free beta subunit) gave a signal in this assay (See Figure 7). There were no lower molecular weight fragments identified by the B152-B207* assay. The hCG free beta analyte was measured in urine described in Figure 2 and was found to make a negligible contribution to over all hCG immunoreactivity in these specimens.

001050 "STAGE 500

Molecules recognized by monoclonal antibody B152 in urine and pituitary extracts. In order to define the nature of the hCG isoforms recognized by B152, high resolution gel filtration columns of both pituitary
5 extracts and postmenopausal urine concentrates were used. The rationale for use of pituitary extracts is to determine cross-reactive molecules, specifically those which are glycosylated, which are plentiful in pituitary which contains the entire family of glycoprotein
10 hormones, hLH, hTSH, and hFSH as well as free subunits and the pituitary form of hCG. Two peaks are detected in both of these cases. Only one peak was detected in similar studies of pregnancy urine concentrates as described earlier. In the pituitary, it is likely that
15 the larger molecule is pituitary hCG (70K) while the smaller sized molecule is hLH. Since hLH is present at 100x or so as compared to pituitary hCG, the apparent similar concentration of immunoreactivity indicates that B152 has reduced cross-reactivity to hLH as compared to
20 hCG. Likewise, both hCG and hLH occur in postmenopausal urine, again with much more hLH than hCG and the B152 pattern is similar to that of the pituitary extract. These results show that B152 is generally hCG specific except for cross reactivity to hLH (as shown by standard
25 cross-reaction studies in Table I) and that its carbohydrate specificity is both to the protein portion as well as to the carbohydrate moieties of hCG (and to a lesser extent of hLH) since it does not react with the multitude of other glycoylated proteins present in the
30 pituitary nor with those in postmenopausal urine except for hCG or hLH-related molecules.

Serum and urine specimens were analyzed using two
35 assays, B109-B108* and B152-B207*, which recognize the difference in molecular isoforms of hCG. See Table I. The *in vitro* bioassay for hLH/hCG is described above. (See Figure 1). The immunometric assay employs 96-well

0010001512000000

microtiter plate technology. The coating antibody, at a concentration determined to provide the most satisfactory combination of sensitivity and range, is applied to the microtiter wells (Immulon IV, Dynatech Laboratories) in carbonate buffer (0.2M, pH 9.5). The plates are incubated with the coating solution at 4°C, overnight, then aspirated, washed with washing solution (0.05% Tween, 0.15N NaCl), and blocked with a 1% solution of BSA (three hours at room temperature). The BSA solution is aspirated and the appropriate hCG standards (200µL/well), in buffer B (PBS/0.1% bovine IgG/0.1% sodium azide), or in hCG free serum (Chemicon, Inc.), or hCG free urine, as appropriate to the specimen matrix, and specimens are added to the wells. The plates are sealed with plate sealers, and incubated overnight at 4°C. The controls, specimens, and standards are then aspirated, the plates washed 5 times with washing solution, and iodinated detection antibody in buffer B (200 uL/well, 100,000 cpm/well) added and incubated overnight at 4°C. The wells are again aspirated, washed 5 times with washing solution, separated and counted (Packard Cobra gamma counted). Values are interpolated from a smoothed spline transformation of the count data. This assay procedure, as well as assay validation has been previously reported (O'Connor, J. F., et al., 1988).

Creatinine analysis, when urine values are normalized to creatinine, is performed in a microtiter plate format following a modification of the Tausky procedure (Tausky, H. H., 1954).

Descriptive statistical and graphical methods are applied to measures of serum and urine samples from normal healthy pregnancies to identify the distributions a) between patient first trimester average B152 levels, B109 levels and B152/B109 ratio; b) between patient

00T080 " 5120E960

variability in time to B152/B109 ratio reaching 1.00; and c) between patient variability in time to B152/B109 ratio declining by 1/3rd from first trimester maximum levels. The variability in the timing of the crossover
5 in the ratio of these two analytes provides an empirical basis from which to estimate the value of these markers as biochemical signatures of a viable third trimester fetus.

10 Comparison of the assay profile of healthy normal pregnancies to those of unsuccessful pregnancies from failed IVF implantations, two non-parametric hypotheses are available: 1) the proportion of pregnancies in which the B152/B109 ratio falls below 1.00 is no different in
15 healthy normal and unsuccessful IVF pregnancies; 2) the proportion of pregnancies in which the B152/B109 ratio declines by 1/3rd from first trimester maximum levels is no different in healthy normal and unsuccessful IVF pregnancies. These hypotheses can be tested as a
20 difference between two proportions. For example, a comparison of week 14 vs. week 9, week 13 vs. week 6, week 12 vs. week 5 or week 11 vs. week 4 pregnancies to show a reversal of the B152/B109 ratio in healthy normal pregnancies and unsuccessful IVF implantations,
25 respectively. The power analyses apply to an outcome defined as the time at which the B152/B109 ratio declines by 1/3rd from first trimester maximum levels, although this outcome would necessarily provide earlier detection of pregnancy failure than the reversal of the
30 B152/B109 ratio. Patterns of results less discriminantly different from these indicate a rejection of the dichotomous outcome of B152/B109 ratio reversal as a clinically meaningful marker of pregnancy failure.

35 Alternatively, the same two non-parametric hypotheses can be recast as parametric hypotheses by considering the timing of the biochemical events within the assay

096924 090400
007000 570000

profile of healthy normal pregnancies and unsuccessful pregnancies from failed IVF implantations: 1) the time at which the B152/B109 ratio falls below 1.00 is no different in healthy normal and unsuccessful IVF pregnancies; 2) the time at which the B152/B109 ratio declines by 1/3rd from first trimester maximum levels is no different in healthy normal and unsuccessful IVF pregnancies. Of course, the objective is to provide an empirical basis from which clinicians may counsel their patients. Thus, it is important to adopt a logistic model for this component of the data analysis. With pregnancy success as the outcome, logistic models allow the estimation of the (symmetrical) hypothesis of increase in risk of pregnancy failure for each additional week where either the B152/B109 ratio has failed to decline by one third from first trimester baseline maximum values or the B152/B109 ratio has failed to become less than 1.00 (measured in weeks). The logistic model enables specification of the time at which results indicate a particular pregnancy exceeds an a priori defined likelihood of failure, given assay data regularly available during pregnancy, and allows incorporation of other risks for pregnancy failure in the same data analytic framework to assess the relative contribution of threats to pregnancy loss. The Cox proportional hazard model may be used to examine predictors of the crossover rates. Mixed effects models can also analyze repeated measures of the B152/B109 ratios taken during entire cycles. These models are particularly useful since they allow inclusion of incomplete and imbalance data (i.e. data with missing values and unequal timing of data collection), to estimate effects of time-varying covariates, to model dependency structure of repeated measures and to model possible heterogeneity of the ratio measures within each experimental group.

092001 5720E960

B152 hCG isoforms isolated from early pregnancy urine and determination of their protein and carbohydrate structures. Using the already developed scheme of concentration and immunoaffinity extraction of urine, hCG molecules are isolated from urine collected from women in early pregnancy for both protein and carbohydrate analyses. According to one approach, molecules are isolated from HPLC fractions, digested with proteases before and after reduction of disulfide bonds, examination of the resultant peptides by mass spectrometry and/or sequence analysis, isolation of carbohydrate moieties after glycosidase digestions and determination of carbohydrate structures by a combination of specific glycosidases and retention times on specialized anion exchange columns as compared to know branch-chain oligosaccharide standards. In a similar approach, the final purification stage for the isolated hCG isoforms is SDS gel electrophoresis. Both protease digests and glycosidase digests are performed on the blotted and cutout band. This method results in greater purity of the protein and less artifactual errors due to contamination by carbohydrates which are not in the purified protein but are derived from outside contaminants.

Carbohydrate compositional analyses and oligosaccharide branched chain identifications. The MALDI TOF mass spectrometric method may be used to confirm oligosaccharide structures by using specific glycosidases on the glycopeptides and determining the change in molecular weight as the sugars are digested off the glycopeptide. Only the hCG beta COOH peptide can be expected to contain O-linked sugar moieties. These are of special interest since it is thought that B152 has significant reaction with this region. The structures of this region can be determined in a similar fashion using enzymes that specifically release O-linked

007220 "SECRET" 000

on B152 hCG isoforms of early successful pregnancy, hCG from third trimester pregnancy, and the reference urinary hCG preparation, CR 127.

5 Example 3: B152 and B151 immunoreactivity in non-trophoblastic malignancy.

With the exception of trophoblastic disease and testicular cancer, hCG is expressed in the blood of
10 about 20% of patients with all other types of cancer (Hussa, R. O., 1987). HCG beta core fragment in the urine has a significantly higher level of expression, especially in gynecological malignancy. Since the B152 antibody was developed to a form of hCG
15 produced in a malignancy, it was of interest to examine the expression of B152 and nicked hCG immunoreactivity (B151) in non-trophoblastic malignancy. Accordingly, blood and urine derived from men undergoing chemotherapy for prostate cancer or women for ovarian cancer were
20 evaluated for the expression of hCG isoforms in plasma and urine. It is significant that in prostate cancer, B152 hCG immunoreactivity is found in the blood and urine of prostate cancer patients in instances when there is no hCG detected by B109-B108*. In ovarian
25 cancer patients evaluated, there is evidence of nicked hCG in the blood, even in the absence of both B109 and B152 immunoreactivity. Neither of the above groups demonstrated the presence of hCG immunoreactivity when the standard pregnancy derived hCG assay was employed.
30 It is reassuring to find that nicked hCG, the existence of which has been documented by several investigators, can be found and reliably measured in a clinical setting.

35 Experimental Discussion

In the course of these studies, a potentially important new signal was observed in the urine of women early in pregnancy, namely an epitope of a form of hCG which may

001000" ST000950

indicate the likely success of carrying a pregnancy. Likewise, absence of this signal may indicate that EPL will occur. Since EPL can be a very sensitive marker of environmental toxins (Hakim, R. B., et al., 1995) and is frequently used as an epidemiological marker of exposure, the finding of this epitope provides a powerful tool for monitoring the safety of the environment. In addition, this assay facilitates increasing the success rate of IVF infertility programs since the predictive value of the new measuring system would rapidly indicate successful approaches. Described herein is the novel and completely unexpected finding that successful pregnancies display a high content of unique isoforms of hCG that are maintained for the first few weeks of pregnancy and then rapidly decline as pregnancy progresses. Based on properties of the immunoassay system, it is hypothesized that these hCG isoforms may be hyperglycosylated. This is a striking observation never reported nor suspected earlier. Carbohydrate analyses (Elliot, M., 1997) demonstrate that C5 hCG employed as immunogen for antibody B152, contains two times the monoantennary content and three times the tri-antennary content of branch chain sugars as compared to the CR series of natural pregnancy urinary hCG. In addition, the O-linked carbohydrates are mostly tetrasaccharide instead of disaccharide in C5 as compared to CR 127 hCG. (CR 127 hCG is similar to the WHO preparation, the third international hCG standard, which was CR 119 hCG, prepared by Canfield and Birken twenty years ago but still in use today) (Birken, S., et al., 1991a). B152 recognizes C5 hCG much better than nicked CR127 hCG or non-nicked CR 127 hCG (Birken, S., et al., 1993). In addition, JAR cell type hCG is known to contain a similar array of carbohydrate moieties. It was found to be recognized by B152 similar to the early pregnancy isoforms in healthy pregnancies. The observation that the hCG isoform produced by JAR cells

CONFIDENTIAL

in culture (B152/B109 ratio) is similar to that found in early pregnancy hCG isoforms supports the hypothesis that the production of a type of hCG with a particular glycosylation pattern is a prerequisite for a viable pregnancy. This glycosylation pattern is not characteristic of the hCG of later pregnancy.

A variety of pregnancy disorders are testable. One category of patients consists of those women who experience a high rate of recurrent abortions. Even in populations with no known fertility problems, the total rate of pregnancy loss is 32% (EPL plus clinically recognized abortion) (Wilcox, A. J., et al., 1988). The risk of recurrent abortion increases with the number of spontaneous abortions experienced in the past, reaching an incidence of 32% after three consecutive abortions. (Hill, J. A., and Anderson, D. J., 1990). Probable causes of recurrent spontaneous abortion, comprising genetic, infectious, hormonal imbalance, or immunologic factors can be established in less than 60% of all spontaneous abortions, leaving 40+% of spontaneous abortions with a completely unestablished etiology. These facts, taken together with reports establishing that the administration of exogenous hCG can be an effective therapy in subjects with a history of recurrent spontaneous abortion (Quenby, S., and Farquharson, R. G., 1994; Harrison, R. F., 1985) lends support to the hypothesis that a disproportionate production of the ineffective isoforms of hCG in early pregnancy is a causal factor in both early pre-clinical loss as well as in spontaneous abortion.

A second category includes women undergoing embryo transfer. These patients provide several distinct advantages: The patients undergoing this procedure are not treated with crude hCG preparations, making measurement of hCG isoforms easy and decisive since all

007080 "STRE880

hCG forms derive from the embryo none from any injected hCG preparations. Second, is the opportunity to monitor the nature of the isoforms from day 9 of a successful pregnancy. Third, is the ability to obtain large volumes of urine to purify the early pregnancy isoforms to determine their structures. Fourth, since pregnancy loss is from 50% to 70% in this population, the loss can be defined as due to lack of the essential hCG isoform recognized by B152 or due to other causes. Comparison of early pregnancies in populations of women not undergoing *in vitro* fertilization procedures with those undergoing embryo implantation can, thus, assess whether pregnancy loss situations present similar or different patterns of hCG isoforms during the process. The mechanism of pregnancy loss in the general population as compared with the much higher rate of embryo loss in IVF programs may be different. Additionally, it has been established that the hCG produced in choriocarcinoma has differences in carbohydrate structures, sialic acid content and biological activity (Wide, L., and Hobson, B., 1987; Elliot, M., et al., 1997; Husa, R. A., 1987). Since B152-B207* assay incorporate monoclonal antibodies raised against an immunogen derived from choriocarcinoma, specimens may be evaluated from patients with gestational trophoblastic disease in order to determine whether the above assays recognize the hCG produced in these conditions with greater sensitivity and specificity than do assays based on the hCG of normal pregnancy, as is apparently the case for the hCG produced in testicular and ovarian cancer.

There are few reports of changes of carbohydrate content of hCG-related molecules during pregnancy. Blithe and colleagues studied free alpha subunit of hCG whose carbohydrate content differs from that of alpha within hCG by additional carbohydrate antennae and fucose. The carbohydrate of free alpha becomes increasingly complex

001080-5120E950

in terms of more branches and higher content as pregnancy proceeds. It has also been reported that the quantity of fucose increased in both hCG and in free alpha as pregnancy proceeded (Skarulis, M. C., et al., 5 1992). Thus, the literature indicates increasing content and complexity of carbohydrate of hCG and free alpha subunits. However, immunological data using the B152 monoclonal antibody, implies a progression to simpler carbohydrate content during pregnancy. Since 10 the beta COOH-region's O-linked carbohydrates may be involved in the epitope recognized by B152, it is conceivable that the carbohydrate structures of this region may be altered in a different pattern from the N-linked glycans studied by Blithe and colleagues 15 (Skarulis, M. C., et al., 1992; Blithe, D. L., and Iles, R. K., 1995). Data from Skarulis et al. indicate that heterodimeric hCG may contain additional fucose but do not provide data that this late pregnancy hCG becomes hyperglycosylated as does free alpha.

20 Other studies indicated that the forms of hCG during EPL likely differ in biological activity from those hCG isoforms in successful pregnancies (Ho, H.-H., et al., 1997). The *in vitro* bioassays employed in those studies 25 are unsuitable for large-scale studies and are not as reliable as the immunoassays described herein. Furthermore, it is likely that *in vivo* assays may give different results since *in vitro* and *in vivo* assays sometimes give completely disparate results. In this 30 case, *in vivo* and clearance assays are most important in order to identify whether the hCG isoforms are truly more potent in the whole animal and to identify the reasons for the increased potency. Thus *in vitro* and *in vivo* bioactivities of the early pregnancy isoforms of 35 hCG are highly significant.

Carbohydrate differences is a widely accepted

000000 " 572000000

Example 4: Diagnosis of Gestational Trophoblast Disease.

5 An important application of the B152 (early hCG
isoform)/B109 (late hCG isoform) ratio analysis
described herein above is in the very early (and facile)
diagnosis of gestational trophoblast disease. Examples
of gestational trophoblast disease include
choriocarcinoma or hydatidiform mole. In normal
10 pregnancy, the ratio of B152/B109 of the two isoforms of
hCG rapidly decreases, eventually inverting. In
gestational trophoblast disease including
choriocarcinoma or hydatidiform mole, the ratio is
initially higher than found in normal pregnancy, but
15 does not diminish during the course of the apparent
pregnancy. This approach provides a highly sensitive
and specific diagnostic marker for gestational
trophoblast disease.

20 Other pregnancy disorders in which hCG levels are
abnormally high or abnormally low include Down's
syndrome or other aneuploid pregnancies, ectopic
pregnancy, preeclampsia, and intra-uterine growth
retardation. Because the hCG production in these
25 conditions is quantitatively abnormal compared with
normal pregnancy, an altered ratio of the hCG isoforms
identified by B152 (early hCG isoform) and B109 (late
hCG isoform) can be detected.

30 Thus, the dual isoform analysis (B152/B109) further
provides a method for diagnosing pregnancy disorders and
gestational trophoblast disease.

Materials and Methods

35

Hormones

The non-nicked hCG isolated from the CR127 preparation

007020 "51202350

of hCG was used as a standard in both assays (Birken et al. 1993). The pituitary hCG was isolated as described (Birken et al. 1996). C5, a 100% nicked hCG having extra sugars on both N- and O-linked carbohydrate moieties, purified from the urine of a choriocarcinoma patient (Elliott et al. 1997), was supplied by Dr. Laurence Cole (Yale University School of Medicine). Although the C5 immunogen used in the development of B152 antibody was 100% nicked hCG isoform (i.e. had cleavages in the peptide backbone of loop 2 of the b subunit) the antibody did not discriminate nicked from non-nicked hCG (O'Connor et al. 1998).

The same serial dilutions of non-nicked hCG, pituitary hCG and C5 were used for binding characterization in hCG assays. Hormone concentrations of initial stock standards solutions were determined by amino acid analysis.

Immunoradiometric assays (IRMA)

The methodology used in the construction and validation of the ¹²⁵I-B109-B108* assay has been fully described elsewhere (O'Connor et al. 1988). The B152-B207* assay has also been characterized (O'Connor et al. 1998). Both assays were performed with a slight modification of the published procedure: the capture antibody was adsorbed onto the wells of microtiter plates (Immulon IV, Dynatech, Chantilly, VA) by incubating a 5 µg/ml solution (B109-B108* assay) or 25 mg/ml solution (B152-B207* assay) in coating buffer (0.2 M bicarbonate, pH 9.5) overnight at 4 C. The coating antibody solution was aspirated, the plates washed (wash solution: 0.9% NaCl, 0.05% Tween 20) and blocked with a 1% solution of BSA in

35 The first antibody is a capture, the second antibody with an asterisk is an iodinated detection antibody.

011050" ST000000

PBS with 0.1% sodium azide. Following incubation with the BSA solution (minimum 3 hours at room temperature) the blocking solution was removed, the wells again washed with wash solution and 200 ml/well of the appropriate hCG standards were added in phosphate buffer B (PBS with 0.1% bovine gamma globulin and 0.1% sodium azide). After overnight incubation at 4 C, the plates were again aspirated and washed. The 200ml (50,000 cpm-100,000 cpm) of ¹²⁵I- labeled antibody was added to the wells which were again incubated for 24 h at 4 C. The tracer was aspirated, the plates washed with wash solution, the individual wells placed in glass tubes and the radioactivity determined in a Packard Cobra gamma counter. Doses were determined by interpolation from a smoothed spline transformation of the data points.

All samples were stored frozen at -20 C prior to assay. Because extreme values of sample pH may interfere with antibody binding, the urine pH was adjusted with 1.0M Tris (pH 9.0), 50µl/ml urine prior to assay, so that the final pH was in the range of 7-7.4 (O'Connor et al. 1988). Intra-assay variation was 6% for both assays, inter-assay variation was 12% for B109-B108* and 13% for B152-B207* assays. Sensitivity (least detectable dose) defined as +2SD from the zero calibrator, was 1 fmol/ml for the B109-B108* assay and 2.2 fmol/ml for B152-B207* assay.

Patients samples

Urine samples from IVF patients were a gift from Dr. L. Cole. They included spontaneous abortion (n=14, range of gestational age 1.8-4.1 weeks from ET- embryo transfer), ectopic pregnancies (n=7, gestational age 2.3-4 weeks) and normal pregnancy controls (n=65, encompassing the range 0.6 to 5.4 week from ET). Some of the normal pregnancy urine samples throughout the pregnancies were

contains O-linked sugars. It also releases small, non glycopeptides from both subunits (Pollak, S., et al., 1990, Birken, S., et al., 1987; Birken, S., et al., 1986). Next, the main disulfide-linked core of each hCG subunit, is reduced and carboxymethylated, and separated on reverse phase HPLC at pH 5. At this stage, large peptides are isolated, including the glycopeptides. Each separated glycopeptide is redigested with trypsin and re-separated on HPLC at pH 5. These glycopeptides are next employed for two different methods of sugar chain analysis. One method is the approach of releasing the oligosaccharides by enzymatic digestions using PNGase F for the N-linked glycans. The released glycans can be separated from the peptides by ethanol precipitation, desialyated with neuraminidase, and separated directly on a Dionex CarboPac PA-100 column. Oligosaccharide standards are available from Dionex, Oxford Glycosystems and other companies for calibrating column elution times for various glycans (Hardy, M. R., and Townsend, R. R., 1994, Rohrer, J. S., et al., 1993, Weitzhandler, M., et al., 1993; Townsend, R. R., et al., 1989). Confirmation of the released structures is obtained by performing carbohydrate compositional analysis of eluted glycan peaks as well as performing digestions with specific glycosidases and rechromatographing the modified glycan on the Dionex system (Hardy, M. R., and Townsend, R. R., 1994; Rohrer, J. S., et al 1993; Weitzhandler, M., et al., 1993; Townsend, R. R., et al., 1989; Townsend, R. R., et al., 1991; Townsend, R. R., et al., 1989; Hardy, M. R., and Townsend, R. R., 1989; Townsend, R. R., et al., 1988; Hardy, M. R., et al, 1997; Hardy, M. R., and Townsend, R. R., 1988; Dionex, 1997; Spellman, M. W., 1990; Kumarasamy, R., 1990). The newly modified glycan can be observed to elute at the same time as the appropriate standard oligosaccharide and, in addition, the released monosaccharide can frequently be identified as well (Dionex, 1997). Structure determination is

001150 "S" 20050900

facilitated by the use of specific glycosidases for branch chain cleavage as well as for digestion of individual sugars from each of the branch chains. For example, Endo H cleaves high mannose type and hybrid oligosaccharide chains while glycosidase Endo F2 cleaves biantennary complex types and PNAase F cleaves tri and tetra-antennary chains down to the N-Asn bond.

Competitive receptor binding and in vitro bioassay.

Bioassays are performed with recombinant-engineered CHO cells transfected with the human receptor to LH/CG Cells are maintained in Ham's F-12 medium, 4mM Glutamine, 400ug/ml G418 (Gibco), 5% fetal calf serum, 100IU/ml penicillin, 100ug/ml streptomycin. The cells are removed from the flask surface by versene only.

A competitive receptor assay constructed as follows: The receptor binding assay mixture contains 100 ul of the appropriate dilution of serum/urine samples or hCG dilutions for standard curve, 100ul of ¹²⁵-I-hCG (50,000-100,000cpm) in buffer A(PBS/0.1%BSA) and 100ul of CHO cells (2x10⁵ cells in PBS). The mixture is incubated at 37°C with slight shaking followed by centrifugation for 10 minutes at 750xg. The supernatant is aspirated and the cell pellet is counted in gamma-counter.

In vitro bioassay. Transfected CHO cells are seeded (200,000 cells/well) into a 24 well plate in culture medium and incubated for 2-3 days until the cells reach confluence. Non-transfected CHO cells are included to monitor non-specific response. The medium is removed and replaced with medium containing 1 mM isobutylmethylxanthine with appropriate dilutions of tested serum or urine. The plates are incubated at 37°C for two hours. The supernatant is removed, and the wells washed with Hank's balanced salt solution. The intracellular cAMP is extracted with 95% ethanol, which

is diluted 1:5, (or up to 1:40, depending on cAMP content) in assay buffer provided by the cAMP kit (New England Nuclear). cAMP assay is performed according to manufacturer's instructions. Response is normalized to well protein content (BCA protein assay kit, Pierce, Rockford, IL).

In vivo bioassay is determined by the uterine weight assay in immature female mice, following the procedure of Wide and Hobson (Wide, L., and Hobson, B., 1987). The mice are injected subcutaneously with one third of the total dose of gonadotropin on three consecutive days and killed 72 hours after the first injection. Uteri are dissected free from mesentery, fat and oviducts, blotted to remove intrauterine fluid and weighed to the nearest 0.1mg. Five to ten mice are used at each of these dose levels. The hCG standard preparation used is a nicked hCG. This material may be run concurrently with specimens isolated from first and third trimester pregnancy. Sham saline injection may be used as a control. The response signal is the log mouse uterine weight.

Clearance of hCG isoforms. The clearance of hCG is determined in the rat. Blood (200ul/sample) is obtained at 0, 120, 240, 360 and 480 minutes post injection, from an indwelling catheter in an catheterized external jugular vein, following the procedure described by Newman et al. (Newman, C. B., et al., 1985) and Brown and Hedge (Brown, M. R., and Hedge, G. A., 1972). Briefly, adult male Sprague-Dawley rats (Charles River Laboratories, Wilmington MA), wt 175-225g, are given free access to food and water. Rats are handled for acclimatization for one week after arrival, and several days before the hCG infusion, the rats are cannulated under pentobarbital anesthesia. A 21 gauge stainless steel cannula is inserted into the one external jugular

007080 " ST000000

vein. The placement of the catheter allows for the collection of blood from the unrestrained, unstressed rat. After the animals have recuperated from the cannula implacement , an hCG isoform is injected (10
5 $\mu\text{g/ml}$ sterile saline) through the cannulated vein. Blood samples are obtained at the four time intervals listed above. The blood is allowed to clot and the serum separated and stored at -80°C for immunometric assays specific for different hCG isoforms.

10

Clearance rate of the isoforms of hCG from the circulation of the rat are estimated by computer fitting the concentration data to an equation of the general form:

15

Concentration= $Ae^{-\alpha t} + Be^{-\beta t}$ at time t ; A and α are parameters of the rapid component and B and β are parameters of the slow component. The metabolic clearance rate (MCR) is calculated as $\text{MCR} = \text{Dose}/(A/\alpha + B/\beta)$ and the initial volume of distribution is
20 calculated from $V_d = \text{Dose}/(A+B)$. The MCR is normalized to body weight for statistical analysis, which is performed using ANOVA with Duncan's range test for determination of significance (Cassals, J. W., et al., 1989).

25

Mice. The mouse species used in the experiments described herein are Balb/c mice, aged 12-20 weeks old and adult Sprague-Dawley rats of either sex. Mice used for the production of monoclonal antibodies through ascites and for the determination of *in vivo* biological
30 activity as described. Balbc/c mice are used because hybridoma cell lines were developed using Balb/c splenocytes.

References

- Amano, J., R. Nishimura, S. Sato, and A. Kobata. 1990. *Glycobiology*. 1:45-50.
- 5 Armstrong, E. G., P. H. Ehrlich, S. Birken, J. P. Schlatterer, E. Siris, W. C. Hembree, and R. E. Canfield. 1984. *J. Clin. Endocrinol. Metab.* 59:867-874.
- Baenziger, J. U. 1994. *FASEB J.* 8:1019-1025.
- 10 Bahl, O. P., L. Marz, and W. R. Moyle. 1995. pp. 125-44. In Anonymous In: Dufau ML, Means AR, ed. Hormone binding and target cell activation in the testis. New York, Plenum Press, 1974.
- 15 Berger, P., Schwarz, S., Spottl, G., Wick, G. and Mann, K. (1993) Variants of human chorionic gonadotropin from pregnant women and tumor patients recognized by monoclonal antibodies. *J. Clin. Endocrinol. Metab.* 77, 20 347-351.
- Birken, S., M. A. Kolks, S. Amr, B. Nisula, and D. Puett. 1987. *Endocrinology* 121:657-666.
- 25 Birken, S., Gawinowicz, MA, Kardana, A., Cole, LA. 1991. *Endocrinology* 129: 1551-1558.
- Birken, S. , Kovalevskaya, G. , O'Connor, J. *Mol Cell Endocrinol*, 1996, 125:121-131.
- 30 Birken, S., M. A. Gawinowicz Kolks, S. Amr, B. Nisula, and D. Puett. 1986. *J. Biol. Chem.* 261:10719-10727.
- 35 Birken, S., Y. Chen, M. A. Gawinowicz, J. W. Lustbader, S. Pollak, G. Agosto, R. Buck, and J. O'Connor. 1993. *Endocrinology* 133:1390-1397.

027060 "STAGE 50

- Birken, S., Y. Maydelman, M. A. Gawinowicz, A. Pound, Y. Liu, and A. S. Hartree. 1996b. *Endocrinology* 137:1402-1411.
- 5 Blithe, D. L. and R. K. Iles. 1995. *Endocrinology* 136:903-910.
- Braun, J. R., T. E. Willnow, S. Ishibashi, G. Ashwell, and J. Herz. 1996. *J. Biol. Chem.* 271:21160-21166.
- 10 Brown, M. R. and G. A. Hedge. 1972. *Neuroendocrinology.* 9:158-174.
- Browne, E. S., M. V. Flasch, M. R. Sairam, and V. K. Bhalla. 1990. *Biochim. Biophys. Acta* 1033:226-234.
- 15 Cassals, J.W., Mann, K., Blithe, D.L., Nisula, B.C., Wehmann, R.E. 1989. *Cancer* 64:2313-2318.
- 20 Chang, P.L., Canfield, R.E., Ditkoff, E.C., O'Connor, J.F., Sauer, M.V., 1998. *Fertil. Steril.*, 69:412-414.
- Chmielewski, C. 1992. *ANNA. J.* 19:34-38.
- 25 Cole, L.A. and Kardana, A. (1992) Discordant results in human chorionic gonadotropin assays. *Clin. Chem.* 38, 263-270.
- Cole, L. A., A. Kardana, P. Andrade-Gordon, M. A. Gawinowicz, J. C. Morris, E. R. Bergert, J. O'Connor, and S. Birken. 1991a. *Endocrinology* 129:1559-1567.
- 30 Cole, L. A., A. Kardana, F. C. Ying, and S. Birken. 1991b. *Yale J. Biol. Med.* 64:627-637.
- 35 Cole, L. A., A. Kardana, S. Y. Park, and G. D. Braunstein. 1993. *J. Clin. Endocrinol. Metab.*

001050" ST000000

76:704-710.

Cole, L. A., S. Birken, and F. Perini. 1985. *Biochem. Biophys. Res. Commun.* 126:333-339.

5

Diaz-Cueto, L., Barrios-de-Tomasi, J., Timossi, C., Mendez, J.P. and Ulloa-Aguirre, A. (1996) More in-vitro bioactive, shorter-lived human chorionic gonadotrophin charge isoforms increase at the end of the first and during the third trimesters of gestation. *Mol. Hum. Reprod.* 2, 643-650.

10

Dionex. 1997. *Technical Note 42*

15

Elliott, M. M., A. Kardana, J. W. Lustbader, and L. A. Cole. 1997. *Endocrine*, 7:15-32.

Ellish, N.J., Saboda, K, O'Connor, J, Nasca, P.C., Stanek, EF, Boyle, C. *Hum Reprod* 1996 11:406-412.

20

Fares, F. A., N. Sukanuma, K. Nishimori, P. S. LaPolt, A. J. Hsueh, and I. Boime. 1992. *Proc. Natl. Acad. Sci. U. S. A.* 89:4304-4308.

25

Fein, H.G., Rosen, S.W. and Weintraub, B.D. (1980) Increased glycosylation of serum human chorionic gonadotropin and subunits from eutopic and ectopic sources: comparison with placental and urinary forms. *J Clin Endocrinol Metab* 50, 1111-1120.

30

Fiete, D., V. Srivastava, O. Hindsgaul, and J. U. Baenziger. 1991. *Cell* 67:1103-1110.

35

Grotjan, H. R. J. and L. A. Cole. 1989. In *Microheterogeneity of Glycoprotein Hormones*. H. R. J. Grotjan and L. A. Cole, editors. CRC Press, Boca Raton. 219-237.

Hakim, R. B., R. H. Gray, and H. Zacur. 1995. *Am. J. Obstet. Gynecol.* 172:1510-1517.

5 Hardy, M. R. and R. R. Townsend. 1988. *Proc. Natl. Acad. Sci. U. S. A.* 85:3289-3293.

Hardy, M. R., R. R. Townsend, and Y. C. Lee. 1997. *Anal Biochem* 1988 Apr;170(1):54-62,

10 Hardy, M. R. and R. R. Townsend. 1989. *Carbohydr. Res.* 188:1-7.

Hardy, M. R. and R. R. Townsend. 1994. *Methods Enzymol.* 230:208-225.

15 Harrison, R. F. 1985. *Europ. J. Obstet. Gynec. reprod. Biol.* 20:159-168.

20 Hill, J. A. and D. J. Anderson. 1990. *Archsm. Immun. Ther. Exp.* 38:111-119.

Ho, H-H, O'Connor, JF, Overstreet, JW, Lasley, BL. *Early Pregnancy: Biology and Medicine*, 1997, (in press).

25 Hoermann, R., G. Spoetl, M. Grossmann, B. Saller, and K. Mann. 1997. *J. Clin. Invest.* 71:953-960.

30 Hoermann, R., P. Berger, G. Spoetl, F. Gillesberger, A. Kardana, L. A. Cole, and K. Mann. 1994. *Clin. Chem.* 40:2306-2312.

Hussa, R. O. 1987. *The Clinical Marker hCG*. Praeger, New York.

35 Kagimoto, A., R. Sakakibara, N. Fukushima, N. Ikeda, and M. Ishiguro. 1995. *Biol. Pharm. Bull.* 18:810-817.

Kardana, A., G. D. Braunstein, and L. A. Cole. 1996. *Oncol. Res.* 8:13-16.

Kardana, A. and L. A. Cole. 1992. *Clin. Chem.* 38:26-33.

5

Karlsson, F. A., P. Burman, O. Kampe, J. E. Westlin, and L. Wide. 1993. *Acta Endocrinol. (Copenh)* 129:291-295.

Kawasaki, T. and G. Ashwell. 1976. *J. Biol. Chem.* 251:1296-1302.

10

Kovalevskaya, G, Birken, S, O'Connor, J, Schlatterer, J, Maydelman, Y, Canfield, R. *Endocrine*, 1995, 3:881-887.

15

Kumarasamy, R. 1990. *J. Chromatogr.* 512:149-155.

Lasley, B. L., P. Lohstroh, A. Kuo, E. B. Gold, B. Eskenazi, S. J. Samuels, D. R. Stewart, and J. W. Overstreet. 1995. *Am. J. Ind. Med.* 28:771-781.

20

Maack, T., C. H. Park, and M. J. F. Camargo. 1985. In *The kidney: Physiology and pathophysiology*. D. W. Seldin and G. Giebisch, editors. Raven Press, New York. 1773-1803.

25

Matzuk, M. M., A. J. Hsueh, P. Lapolt, A. Tsafiriri, J. L. Keene, and I. Boime. 1990. [published erratum appears in *Endocrinology* 1990 Apr;126(4):2204]. *Endocrinology* 126:376-383.

30

Moyle, W. R., O. P. Bahl, and L. Marz. 1975. *Journal of Biological Chemistry* 250:9163-9169.

35

Newman, C. B., S. L. Wardlaw, and A. G. Frantz. 1985. *Life Sci.* 36:1661-1668.

Nishimura, R., T. Kitajima, K. Hasegawa, K. Takeuchi,

and M. Mochizuki. 1989. *Jpn. J. Cancer Res.* 80:968-974.

Nishimura, R., T. Utsunomiya, K. Ide, T. Kitajima, H. C. Chen, J. L. Strobel, R. O. Husa, and M. Mochizuki.
5 1987. *Jpn. J. Cancer Res.* 78:833-839.

Nishimura, R., K. Ide, T. Utsunomiya, T. Kitajima, Y. Yuki, and M. Mochizuki. 1988. *Endocrinology* 123:420-425.

10 O'Connor, J., G. Kovalevskaya, S. Birken, J. P. Schlatterer, D. Schechter, D. McMahon, and R. E. Canfield. 1998. *Hum. Reprod.* 13:826-835.

O'Connor, J. F., S. Birken, J. W. Lustbader, A.
15 Krichevsky, Y. Chen, and R. E. Canfield. 1994. *Endocr. Rev.* 15:650-683.

O'Connor, J. F., J. P. Schlatterer, S. Birken, A. Krichevsky, E. G. Armstrong, D. McMahon, and R. E. Canfield. 1988. *Cancer Res.* 48:1361-1366.
20

Odell, W. D. and J. Griffin. 1989. *J. Clin. Endocrinol. Metab.* 69:528-532.

25 Odell, W. D. and J. Griffin. 1987. *N. Engl. J. Med.* 317:1688-1691.

Pollak, S., S. Halpine, B. T. Chait, and S. Birken. 1990. *Endocrinology* 126:199-208.

30 Puisieux, A., D. Bellet, F. Troalen, A. Razafindratsita, C. Lhomme, C. Bohuon, and J. M. Bidart. 1990. *Endocrinology* 126:687-694.

35 Quadri, K. H., J. Bernardini, A. Greenberg, S. Laifer, A. Syed, and J. L. Holley. 1994. *Am. J. Kidney Dis.* 24:416-420.

001030"5720360

- Quenby, S. and R. G. Farquharson. 1994. *Fertil. Steril.* 62:708-710.
- 5 Ravindranath, N., N. S. Srilatha, M. R. Sairam, and N. R. Moudgal. 1992. *Indian J. Exp. Biol.* 30:982-986.
- Rohrer, J. S., G. A. Cooper, and R. R. Townsend. 1993. *Anal. Biochem.* 212:7-16.
- 10 Sairam, M. R. and L. G. Jiang. 1992. *Mol. Cell Endocrinol.* 85:227-235.
- Sairam, M. R. 1989. *FASEB J.* 3:1915-1926.
- 15 Sairam, M. R., J. Linggen, and G. N. Bhargavi. 1988. *Biosci. Rep.* 8:271-278.
- Skarulis, M. C., R. E. Wehmann, B. C. Nisula, and D. L. Blithe. 1992. *J. Clin. Endocrinol. Metab.* 75:91-96.
- 20 Spellman, M. W. 1990. *Anal. Chem.* 62:1714-1722.
- Stanton, P. G., G. Poznek, P. G. Burgon, D. M. Robertson, and M. T. W. Hearn. 1993. *J Endocrinol.* 25 138:529-543.
- Szkudlinski, M. W., N. R. Thotakura, I. Bucci, L. R. Joshi, A. Tsai, J. East-Palmer, J. Shiloach, and B. D. Weintraub. 1993. *Endocrinology* 133:1490-1503.
- 30 Szkudlinski, M. W., N. R. Thotakura, J. E. Tropea, M. Grossmann, and B. D. Weintraub. 1995. *Endocrinology* 136:3325-3330.
- 35 Taussky, H. H. 1954. *J. Biol. Chem.* 208:853-861.
- Thotakura, N. R., M. W. Szkudlinski, and B. D.

What is claimed is:

1. A method of predicting the likelihood of a negative pregnancy outcome in a female subject comprising:

- 5 (a) contacting a sample from the subject with a capture antibody which specifically binds to an early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG;
- 10 (b) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample; and
- 15 (c) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (b) with the amount determined for temporally matched, normal pregnant subject, wherein the relative absence of the early pregnancy associated molecular isoform of hCG in the sample indicates a negative outcome of pregnancy for the subject.
- 20
- 25

2. A method of predicting the likelihood of a negative pregnancy outcome in a female subject comprising:

- 30 (a) contacting a sample from the subject with a capture antibody which specifically binds to an early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG;
- 35 (b) contacting any complex formed in step (a) with a labelled detection antibody under

001020 " 5120E950

conditions permitting binding to the complex the capture antibody and the hCG isoform;

5 (c) measuring the amount of labeled detection antibody bound to the complex so as to thereby determine the amount of the early pregnancy associated molecular isoform of hCG in the sample; and

10 (d) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (b) with the amount determined for a normal pregnant subject, wherein the relative absence of the early pregnancy associated molecular isoform of hCG in the sample indicates a negative outcome of pregnancy for the subject.

15 3. The method of claim 1, step (a) further comprising a detection antibody which specifically binds to hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG.

20 4. The method of claim 3, wherein the capture antibody is B152.

25 5. The method of claim 4, wherein the detection antibody is B207.

30 6. A method of predicting the likelihood of a negative pregnancy outcome in a female subject comprising:

35 (a) contacting a sample from the subject with a capture antibody which specifically binds to an early pregnancy associated molecular isoform of hCG under conditions permitting formation of a

001080" 5120E960

molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG.

5 9. The method of claim 1, wherein the sample is a urinary sample or a blood sample.

10. The method of claim 1, wherein the sample is an aggregate sample taken from at least one day.

10

11. The method of claim 1, wherein the antibody is labeled with a detectable marker.

15

12. The method of claim 11, wherein the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin.

15

13. The method of claim 12, wherein the radioactive isotope is I¹²⁵.

20

14. A method of predicting pregnancy outcome in a subject by determining the amount of an early pregnancy associated molecular isoform of hCG in a sample comprising:

25

(a) contacting a capturing antibody which specifically binds to the early pregnancy associated molecular isoform of hCG with a solid matrix under conditions permitting binding of the antibody with the solid matrix;

30

(b) contacting the bound matrix with the sample under conditions permitting binding of the early pregnancy associated molecular isoform of hCG present in the sample with the capturing antibody;

35

(c) separating the bound matrix and the sample;

(d) contacting the separated bound matrix with a detecting antibody which specifically binds to

027020-5722950

hCG under conditions permitting binding of antibody and antigen in the sample;

5 (e) measuring the amount of bound antibody on the bound matrix, thereby determining the amount of early pregnancy associated molecular isoform of hCG in the sample;

10 (f) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (e) with the amount determined for temporally matched, normal pregnant subject(s), wherein amounts of the early pregnancy associated molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG in temporally matched pregnant samples indicates a positive outcome, and a less amount indicates a negative outcome of pregnancy for the subject.

15 15. The method of claim 14, further comprising:

20 (a) removing of the sample from the matrix; and
(b) washing the bound matrix with an appropriate buffer.

25 16. The method of claim 14, wherein the capturing antibody is B152.

17. The method of claim 14, wherein the detecting antibody is B207.

30 18. The method of claim 14, step (a) further comprising a second capturing antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and
35 the early pregnancy associated molecular isoform of hCG.

001050-5120E550

isotope is I¹²⁵.

27. A method for determining the amount of early pregnancy associated molecular isoforms in a sample comprising:

- (a) contacting the sample with an antibody which specifically binds to an early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; and
- (b) determining the amount of complexes formed thereby determining the amount of early pregnancy associated molecular isoform of hCG in the sample.

28. The method of claim 27, wherein the antibody specifically binds a region of the early pregnancy associated molecular isoform of hCG comprising a carbohydrate moiety.

29. The method of claim 27, wherein the antibody is produced by the hybridoma cell line accorded ATCC Accession No.HB-12467.

30. The method of claim 27, wherein the antibody is B152.

31. A diagnostic kit for determining the amount of early pregnancy associated hCG is a sample comprising:

- (a) An antibody which specifically binds to an early pregnancy associated molecular isoform; and
- (b) a solid matrix to which the antibody is bound; and
- (c) reagents permitting the formation of a

molecular isoform detected in step (b) and a relative absence of the early pregnancy associated molecular isoform of hCG detected in step (c) indicates the presence of non-trophoblast malignancy in the sample.

5

45. The method of claim 44, wherein the antibody is B152.

10

46. The method of claim 44, wherein the complex formed in step (a) is detected by a detection antibody.

15

47. The method of claim 46, wherein the detection antibody is B207.

48. The method of claim 44, wherein the second antibody is B109.

20

49. The method of claim 44, wherein the complex formed in step (b) is detected by a detection antibody.

25

50. The method of claim 49, wherein the detection antibody is B108.

30

51. The method of claim 44, wherein the non-trophoblast malignancy is ovarian malignancy or prostate malignancy or some other non-trophoblast malignancy.

52. The method of claim 44, wherein the sample is a urinary sample or a blood sample.

35

53. A method for detecting gestational trophoblast disease in a sample from a subject comprising:
(a) contacting a sample with an antibody which

specifically binds to the early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG;

5

(b) contacting the sample with a second antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG;

10

(c) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample due to binding with the first antibody, and late pregnancy associated molecular isoform of hCG in the sample due to binding with the second antibody;

15

(d) determining the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the subject; and

20

(e) comparing the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) over time, wherein a continuing high ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) indicates the presence of gestational trophoblast disease in the subject.

25

30

35

54. The method of claim 48, wherein the antibody is


001000005120E960

B152.


55. The method of claim 48, wherein the second antibody is B109.
- 5
56. The method of claim 48, wherein the gestational trophoblast disease is choriocarcinoma or hydatidiform mole.
- 10 57. The method of claim 48, wherein the sample is a urinary sample or a blood sample.



001080" 3720E 360

FIG. 1




007080 51202.950





- choriocarcinoma hCG (C5)
- pregnancy hCG
- ▲ pituitary hCG

FIG. 4



COTTON SEED

FIG. 5





FIG. 6



007030" 57202350

FIG. 7A





FIG. 7B



- B109-B108* ▲ B152-B207* ◆ B210-B108*
- B151-B207* ▼ B201-C104*

007080" 520E960

DECLARATION AND POWER OF ATTORNEY

As a below-named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHODS FOR PREDICTING PREGNANCY OUTCOME IN A SUBJECT BY hCG ASSAY

the specification of which:
(check one)

X is attached hereto.
was filed on _____ as
Application Serial No. _____
and was amended _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119 (a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International Application which designated at least one country other than the United States, listed below. I have also identified below any foreign application for patent or inventor's certificate, or PCT International Application having a filing date before that of the earliest application from which priority is claimed:

Table with 5 columns: Prior Foreign Application(s) Number, Country, Filing Date, Priority Claimed Yes, Priority Claimed No. Row 1: PCT/US99/02289, PCT, 3 February 1999, X, .

007050 "ST 200 550

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

<u>Provisional Application No.</u>	<u>Filing Date</u>	<u>Status</u>
N/A		

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States Application(s), or Section 365(c) of any PCT International Application(s) designating the United States listed below. Insofar as this application discloses and claims subject matter in addition to that disclosed in any such prior Application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56, which became available between the filing date(s) of such prior Application(s) and the national or PCT international filing date of this application:

<u>Application Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
09/017,976	3 February 1998	Pending
PCT/US99/02289	3 February 1999	Pending

And I hereby appoint

John P. White (Reg. No. 28,678); Christopher C. Dunham (Reg. No. 22,031); Norman H. Zivin (Reg. No. 25,385); Jay H. Maioli (Reg. No. 27,213); William E. Pelton (Reg. No. 25,702); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Wendy E. Miller (Reg. No. 35,615); Richard S. Milner (Reg. No. 33,970); Robert T. Maldonado (Reg. 38,232); Paul Teng (40,837); Richard F. Jaworski (Reg. No. 33,515); Elizabeth M. Wieckowski (Reg. No. 42,226); Pedro C. Fernandez (Reg. No. 41,741); Gary J. Gershik (Reg. No. 39,992); Jane M. Love (Reg. No. 42,812); Spencer H. Schneider (Reg. No. 45,923) and Raymond A. Diperna (Reg. No. 44,063).

and each of them, all c/o Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, New York 10036, my attorneys, each with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to receive the patent, to transact all business in the Patent and Trademark Office connected therewith and to file any International Applications which are based thereon under the provisions of the Patent Cooperation Treaty.

09/017,976; 02/03/99

Please address all communications, and direct all telephone calls, regarding this application to:

John P. White Reg. No. 28,678
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
Tel. (212) 278-0400

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first joint inventor John F. O'Connor

Inventor's signature

Citizenship U.S.A. Date of signature

Residence 35 Clinton Place, New Rochelle, New York 10801

Post Office Address Same as Residence

Full name of joint inventor (if any) Galina I. Kovalevskaya

Inventor's signature

Citizenship Russia Date of signature

Residence 850 West 176th Street, Apt. #5G, New York, New York 10033

Post Office Address Same as Residence

Full name of joint inventor (if any) Steven Birken

Inventor's signature

Citizenship U.S.A. Date of signature

Residence 120 Oneida Avenue, Dumont, New Jersey 07628

Post Office Address Same as Residence

007000 "ST 202.850